



RESPONSE OF INTEGRATED NUTRIENT MANAGEMENT ON THE GROWTH, YIELD AND QUALITY OF *KHARIF* ONION (*ALLIUM CEPA* L.)

S. Krishnaprabu

Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalainagar-608 002, Tamilnadu, India.

Abstract

The field experiment of study the “Response of fertility levels on growth, yield and quality of *Kharif* onion” was conducted at farmers field at Trichy. The experiment comprising 8 fertility levels (Control, 75%, 100% and 125%) recommended dose of NPK vermicompost 2.5 t ha⁻¹ with 25 per cent, 50 per cent and 75 per cent recommended dose of NPK. The application of 50 per cent recommended dose of NPK + vermicompost 2.5 t ha⁻¹ significantly improved the plant height, number of leaves at harvest, equatorial diameter thickness of scale, volume of bulb, yield and TSS, vitamin “C” and allyl propyl disulphide content.

Key words : Recommended dose of NPK, Vermicompost, Yield, Yield attributes quality.

Introduction

Onion is cultivated in rabi season but early *Kharif* and late *kharif* crops are also taken in various parts of India, During October-November there is shortage of onion in the market, which lends to heavy parts of India. During October – November there is shortage of onion in the market, which lends to heavy prices. Therefore, production of onion in *Kharif* season is more important for continuous supply of onion round the years. Integrated nutrient supply approach is not only the liable way for obtaining fairly high productivity with substainal fertilizer economy but a concept of ecological soundness leading to sustainable agriculture (Swaminathan, 1988). Use of vermicompost in conjunction with chemical fertilizer has been found to be promising not only in maintaining and sustaining higher productivity but also providing stability in crop production. The information on the balanced use of chemical fertilizers along with vermicompost for *Kharif* season onion in the state in very scare. Hence, an experiment was conducted to determine best level of inorganic and organic fertilizer for enhancing the growth, yield and quality of *Kharif* onion.

Materials and Methods

An experiment was conducted during *Kharif* seasons of 2018 and 2019 at farmers field Trichy. In split plot design with four replication. To obtain good quality of onion sets they are prepared at there and variety N-53 were taken. The seeds were sown in well prepared nursery bed in the first week of February at closer spacing. Then sets were lifted in the first week of May from the nursery beds and tops were removed and graded. The fertility treatment were like as: control, 75 per cent, 100 per cent and 125 per cent recommended dose of NPK, Vermicompost dose of NPK. The recommended dose of NPK was 100 kg N, 50 kg P₂O₅ and 100 kg Murate Of Potash and calculated according to treatment wise. The full dose of phosphorus and potash and half dose of nitrogen were given at the time of transplanting of sets, and remaining half dose of nitrogen was given 30 days after transplanting and vermicompost was supplied before the planting of sets. The sets were planted 45 x 10 cm distance and both sides of ridges in Aug. month. All the observation were taken and harvesting was determined with the help of hand refractometer at the time of harvesting of bulb. The vitamin “C” content of bulb was determined by diluting the known volume of

*Author for correspondence : E-mail : prabu1977krishna@gmail.com

onion juice with 3 per cent meta-phosphoric acid and titrating with 2, 6 dichloro phenyl indophenols solution (A.O.A.C., 1960), till the faint pink colour was obtained. Allyl propyl disulphide content in onion bulb was determined as pyruvic acid (Hort and Fisher, 1971).

Results and Discussion

It is revealed from the data table 1 that the plant height, number of leaves plant⁻¹ at harvest of onion were significantly affected with different fertility levels. The maximum plant height (5.11 cm), with maximum number of leaves (11.91) were obtain with 50 per cent recommended dose of NPK along with vermicompost 2.51 ha⁻¹ which was at par with 75 per cent recommended dose NPK along with vermicompost 2.5 t ha⁻¹. The improvement in plant height, number of leaves with the application of vermicompost might be due to better moisture holding capacity and supply and availability of major and micro nutrient due to favourable soil condition (Reddy *et al.*, 1998). In case of vermicompost the earthworm carts helps in improvement the soil fertility and availability and nutrients besides some growth stimulating substances excreted by earthworm into their carts. (Senapati *et al.*, 1985).

The yield attributes of onion was significantly influenced with different fertility level. The maximum equatorial diameter (5.62 cm), thickness of scales (0.274 cm), volume of bulb (56.03 cc) and yield (224.29 q ha⁻¹) were recorded with 50 per cent recommended dose of NPK along with vermicompost 2.51 ha⁻¹. However neck

length (6.40 cm) and fresh weight of bulb (95.14 g) were maximum in 75 per cent recommended dose NPK plus vermicompost 2.5 t ha⁻¹ but it was at par with (T₇). The present trend of increase in bulb yield and yield attributes with 5 per cent recommended dose of NPK along with vermicompost 2.5 t ha⁻¹ is in close conformity with findings of Mahendran and Kumar (1997) and Patil *et al.* (2002), Application of organic manure in the form of vermicompost increased the yield attributes and yield. Such increase may be due to release of macro and micronutrient, during the course of microbial decomposition (Singh and Ram, 1982). The beneficial response of FYM vermicompost to yield might also be attributed to availability of sufficient amount of plant nutrients throughout the growth period of crop uptake, plant vigour and yield (Brar and Pasricha, 1998).

The data presented in Table 1 revealed that application of 75 per cent recommended dose of NPK along with vermicompost 2.5 t ha⁻¹ affected the TSS, (12.16), Allyl propyl disulphide (7.23) and vitamin "C" (9.68mg/100 g) content but it was at par with 50 per cent recommended dose of NPK plus vermicompost 2.5 t ha⁻¹. An increased in NPK significantly vitamin "C" content, which was due to helped in vigorous vegetative growth and imparted green colour to foliage which favoured photosynthetic activity of plant so there was greater accumulation of food material. The similar results have been also been reported by Singh *et al.*, (1989); Thabet *et al.*, (1994) and Singh *et al.*, (1996).

Table 1: Response of integrated nutrient management on growth, yield and quality of *Kharif* onion.

Treatments	Plant height (cm)	Number of leaves per plant at harvest	Neck thickness (cm)	Equatorial diameter (cm)	Thickness of scale (cm)	Fresh weight of bulb (g)	Volume of bulb (cc)	Yield (q per ha)	T.S.S. (%)	Allyl-propyl disulphide	Vitamin C (mg/100g)
Control	3842	6.61	5.73	4.07	0.188	56.59	41.63	128.76	12.04	6.57	9.20
75% recommended dose of NPK	52.29	9.80	5.81	4.80	0.212	69.27	48.05	158.51	12.09	6.67	9.23
100% recommended dose of NPK	52.96	10.67	5.94	5.34	0.248	79.53	53.74	182.03	12.15	6.95	9.58
125% recommended dose of NPK	52.82	10.36	6.08	5.25	0.229	76.39	53.50	174.97	12.20	6.81	9.44
Vermicompost 2.51 ha ⁻¹	52.47	10.08	6.21	5.15	0.221	71.11	51.18	161.27	12.22	6.74	9.36
25% recommended dose of NPK + vermicompost 2.5 t ha ⁻¹	52.99	11.17	6.28	5.43	0.263	86.48	54.37	198.10	12.27	7.01	9.62
50% recommended dose of NPK + vermicompost 2.51 ha ⁻¹	53.11	11.91	6.32	5.62	0.274	94.78	56.03	224.29	12.32	7.23	9.66
75% recommended dose of NPK + vermicompost 2.5 t ha ⁻¹	53.16	11.63	6.40	5.60	0.272	95.14	55.15	214.54	12.36	7.07	9.68
S.E.±	0.654	0.121	0.067	0.054	0.0024	0.439	0.664	2.605	0.083	0.067	0.067
C.D. (P=0.05)	1.832	0.338	0.187	0.153	0.0066	1.230	1.860	7.301	0.232	0.187	0.188

References

- A.O.A.C. (1960). Official methods of analysis Washington D.C. 9th Edn., 15-16.
- Brar, B.S. and M.S. Pasricha (1998). Long term studies on integrated use of organic and inorganic fertilizer in maize-wheat- cowpea cropping system on alluvial soil of Punjab. In Long Term Soil Fertilizer Management Through Integrated Plant Nutrient Supply System 14-168, IISS-Bhopal.
- Hort, F.L. and H.J. Fisher (1971). Determination of pyruvic acid in dehydrated onion. In Modern Food Analysis Springer verlog. Berlin, Neidelberg, New Youk, 433-434.
- Mahendran, P.B. and N. Kumar (1997). Effect of organic manures on cabbage cv. HERO. *South Indian Hort.*, **45(5-6)**: 240-43.
- Patil, M.P., M.B. Madalageri and R. Mulge (2002). Fertilizer use economy in onion. International conference on vegetables. Nov. 11 - 14th, 2002. Bangalore *vegetables of sustainable food and nutritional security in the New Millennium*, 215.
- Reddy, R., M.A. Reddy Narayana and Y.T. Reddy Narayana (1998). Effect of organic and inorganic sources of NPK on growth and yield of pea. *Legume Res.*, **21(1)**: 57-60.
- Senapati, B.K., S.C. Pani and A. Kabi (1985). Effect of earthworm and green manuring on paddy production in pot culture. In: National Seminar on *Current Trends in Soil Biology*, HAU, India, 71-75.
- Singh, H., S. Singh, V. Singh, H. Singh, S. Singh and V. Singh (1996). Response of onion to nitrogen and sulphur. *Ann. Agric. Res.*, **17**: 441-444.
- Singh, R.S. and H. Ram (1982). Effect of organic matter on the transformation of inorganic phosphorus in soils. *J. Indian Soc. Soil Sci.*, **30**: 185-189.
- Singh, T., S.B. Singh and B.N. Singh (1989). Effect of nitrogen, potassium and green manuring on growth and yield of rainy season onion. *Narendra Dev. J. Agril. Res.*, **4** : 57-60.
- Thabet, E.M.A., A.A.G. Adballah and A.R.A.G. Mohamed (1994). Productivity of onion grown in reclaimed sandy soil using tafla as affected by water tegimes and nitrogen levels. *Ann. Agric. Sci. Cairo.*, **39**: 337-344.