



Plant Archives

Journal homepage: <http://www.plantarchives.org>

DOI Url : <https://doi.org/10.51470/PLANTARCHIVES.2025.v25.supplement-2.098>

EFFECT OF DIFFERENT GROWING MEDIA AND PLANTING DATES ON GROWTH AND FLOWERING OF CALENDULA (*CALENDULA OFFICINALIS* L.)

Nikita Maurya¹, Ajay Kumar Singh^{1*}, Amit Kanawjia¹, Arbind Kumar Gupta², Vaishali Gangwar³, Balaji Vikram⁴, Ajay Kumar¹, and Amit Kumar Maurya⁵

¹ Department of Floriculture and Landscape Architecture, BUAT, Banda, U.P., India

² Department of Natural Resource Management, BUAT, Banda, U.P., India

³ Department of Crop Physiology, BUAT, Banda, U.P., India

⁴ Department of Post- Harvest Technology, BUAT, Banda, U.P., India

⁵ Department of Horticulture, Institute of Agricultural Sciences, B.H.U., Varanasi, U.P., India

*Corresponding author E-mail: ajayorchid@gmail.com

(Date of Receiving : 28-04-2025; Date of Acceptance : 03-07-2025)

ABSTRACT

The present study was conducted to evaluate the effect of different growing media and planting dates on the vegetative growth and flowering performance of calendula (*Calendula officinalis* L.). The experiment included six growing media combinations and two planting dates: the second fortnight of October and the second fortnight of November. Results revealed that both growing media and planting dates significantly influenced key vegetative and floral traits. The combination of Cocopeat: Sand: FYM: Vermicompost (2:1:1:1) planted in the second fortnight of October (G₄P₁) produced the best results, including maximum plant height, plant spread, number of primary branches, number of flowers per plant, and flowering duration, along with the earliest flowering whereas G₆P₁ Sphagnum moss: Sand: FYM: Vermicompost (2:1:1:1) planted in the second fortnight of October gives maximum flower size. The superior performance under this treatment (G₄P₁) was attributed to improved aeration, nutrient availability, and favorable climatic conditions during early plant development. The findings suggest that the use of cocopeat-based media with early October planting is optimal for enhancing the growth and floral yield of calendula under Bundelkhand conditions.

Keywords: Growing media, planting dates, vegetative growth, flowering performance, calendula (*Calendula officinalis* L.).

Introduction

Calendula (*Calendula officinalis* L.), commonly known as pot marigold, belongs to the family Asteraceae and is one of the most important annual ornamental and medicinal herbs cultivated worldwide. Native to southern Europe and the Mediterranean region. It has gained significant attention not only for its aesthetic appeal in landscapes and home gardens but also for its wide array of therapeutic and industrial uses. However, its growth, flowering behavior, and yield potential are significantly influenced by several factors, particularly planting time and growing media composition. Potted ornamental plants have become increasingly popular not only for home gardens but also for commercial purposes such as weddings,

exhibitions, hotel décor, and restaurant aesthetics. In this context, the demand for lightweight and sustainable growing media has grown significantly, as they facilitate easy handling, transportation, and reusability, making them highly suitable for rental-based ornamental plant businesses (Dubey *et al.*, 2012). The quality of potted plant production is greatly influenced by the type of growing media used, as it plays a vital role in supplying nutrients, supporting root development, and providing mechanical stability to plants in containers (Vendrame *et al.*, 2005). Commonly used substrates such as sand, farmyard manure (FYM), cocopeat, and sphagnum moss not only vary in their physical and chemical properties but also significantly influence plant growth and nutrient uptake (Bechmann-Cavalcante *et al.*, 2009).

The vegetative growth and overall quality of calendula can be significantly enhanced by selecting an appropriate planting time, which also helps to fulfil market and consumer demands (Muhammad *et al.*, 2001). However, the optimal planting period may differ across regions due to variations in environmental factors such as photoperiod, temperature, and light intensity. The growth and flowering response in calendula are primarily influenced by these climatic factors, particularly day length and ambient temperature. To ensure a consistent supply of flowers throughout the year, staggered planting schedules are often adopted. In this regard, the present investigation was undertaken to study the effect of different growing media and planting dates on the growth, flowering, and yield of Calendula, with an objective to identify suitable agronomic practices that can improve its productivity and adaptability under varying environmental conditions.

Materials and Methods

The present investigation was carried out during the year at the Nursery of the Department of Floriculture and Landscape Architecture, Banda University of Agriculture and Technology, Banda. The site is located in the semi-arid agro-climatic zone of Bundelkhand, characterized by low rainfall, high temperature fluctuations, and prolonged dry spells. The soil of the experimental field was sandy loam in texture, low in organic carbon, and medium in available NPK. The experimental crop used was Calendula (*Calendula officinalis* L.), grown from healthy and certified seeds of a commonly procured from a reliable commercial source. The experiment was laid out in a Factorial Completely Randomized Design (FCRD) with three replications, comprising two planting dates as factors and six different growing media compositions, making a total of 12 treatment combinations with total of 108 pots.

Table 1 : Treatment combinations

No.	Notation	Details
T ₁	G ₁ P ₁	Soil + Sand + FYM (2:1:1, v/v) and Planting on 2 nd fortnight of October
T ₂	G ₁ P ₂	Soil + Sand + FYM (2:1:1, v/v) and Planting on 2 nd fortnight of November
T ₃	G ₂ P ₁	Soil + Sand + FYM +Vermicompost (2:1:1:1, v/v) and Planting on 2 nd fortnight of October
T ₄	G ₂ P ₂	Soil + Sand + FYM +Vermicompost (2:1:1:1, v/v) and Planting on 2 nd fortnight of November
T ₅	G ₃ P ₁	Cocopeat + Sand + FYM (2:1:1, v/v) and Planting on 2 nd fortnight of October
T ₆	G ₃ P ₂	Cocopeat + Sand + FYM (2:1:1, v/v) and Planting on 2 nd fortnight of November
T ₇	G ₄ P ₁	Cocopeat + Sand + FYM + Vermicompost (2:1:1:1, v/v) and Planting on 2 nd fortnight of October
T ₈	G ₄ P ₂	Cocopeat + Sand + FYM + Vermicompost (2:1:1:1, v/v) and Planting on 2 nd fortnight of November
T ₉	G ₅ P ₁	Sphagnum peat moss + Sand + FYM (2:1:1, v/v) and Planting on 2 nd fortnight of October
T ₁₀	G ₅ P ₂	Sphagnum peat moss + Sand + FYM (2:1:1, v/v) and Planting on 2 nd fortnight of November
T ₁₁	G ₆ P ₁	Sphagnum peatmoss + Sand + FYM + Vermicompost (2:1:1:1, v/v) Planting on 2 nd fortnight of October
T ₁₂	G ₆ P ₂	Sphagnum peatmoss + Sand + FYM + Vermicompost (2:1:1:1, v/v) Planting on 2 nd fortnight of November

Sowing was done manually on the respective planting dates, with healthy seedling. Regular irrigation and standard cultural practices such as weeding and pest management were followed uniformly for all treatments. Observations were recorded on growth, flowering at proper stages.

Result and Discussion

The appropriate choice of growing medium plays a crucial role in ensuring adequate root anchorage, efficient nutrient and water supply, and proper gaseous exchange between the root zone and the external environment (Abad *et al.*, 2001). Growing substrates that are lightweight, nutrient-rich, and possess good drainage capacity are regarded as highly suitable for promoting optimal growth and development in

flowering plant species (Dubey *et al.*, 2013). The result of the present investigation is given in Tab 2 and Tab 3. Which revealed that the highest plant height was recorded in treatment G₄P₁, which consisted of Cocopeat: Sand: FYM: Vermicompost (2:1:1:1) planted during the second fortnight of October, with a height of 14.44 cm. Conversely, the lowest plant height was observed in treatment G₁P₂, comprising Soil: Sand: FYM (2:1:1) planted in the second fortnight of November, with a value of 10.88 cm. The superior performance of G₄P₁ can be attributed to the favorable physical and biological properties of cocopeat. Its lightweight and porous nature facilitates improved aeration and moisture retention in the root zone, thereby enhancing plant growth. These findings are in agreement with those of Thumar *et al.* (2020), who

reported maximum plant height in a growing medium composed of Soil + Cocopeat + Leaf Mould (2:1:1). The effectiveness of cocopeat and leaf mould is likely due to their enhanced nutrient content and their ability to improve the physicochemical environment of the root zone. Supporting evidence is also provided by Younis *et al.* (2015), who demonstrated the significant role of cocopeat and leaf mould in promoting increased plant height compared to other growing media. With respect to October planting attained significantly greater height. The enhanced vegetative growth under October can be ascribed to more favorable environmental conditions prevailing during early plant establishment, such as optimal temperatures and sufficient light intensity. These conditions likely promote higher metabolic activity, leading to more vigorous growth. Similar observations were reported by Verma *et al.* (2001) and Lakshmi *et al.* (2014) in African marigold cv. Pusa Narangi, where earlier planting dates resulted in increased plant height due to conducive climatic factors. Among the different growing media Tab 2. A significant interaction effect was observed between growing media and planting dates on plant spread in both the East–West (E–W) and North–South (N–S) directions, indicating a differential response to the combined influence of these factors. In the E–W direction, the maximum value of 24.45 cm was recorded in treatment G₄P₁, which comprised Cocopeat: Sand: FYM: Vermicompost (2:1:1:1) and was planted in the second fortnight of October. Similarly, for plant spread in the N–S direction, the highest value of 24.41 cm was also observed in the same treatment (G₄P₁), highlighting the superior horizontal growth achieved under this media and planting date combination. In contrast, the minimum plant spread of 19.96 cm (N–S) was recorded in G₁P₂, which consisted of Soil: Sand: FYM (2:1:1) planted during the second fortnight of November. The enhanced plant spread under the G₄P₁ treatment can be attributed to the balanced composition of the growing media, particularly the inclusion of vermicompost. Vermicompost is rich in organic matter and beneficial microorganisms, which enhance soil structure, nutrient availability, and root development, thereby promoting better canopy expansion. These findings are supported by the results of Nair and

Bharathi (2015) and Deogade *et al.* (2020), who also reported improved plant spread with the use of vermicompost-enriched media. The superiority of October planting may be due to the availability of more favorable environmental conditions—such as moderate temperatures and better light intensity during the early vegetative phase, which likely contributed to enhanced lateral growth. These observations are in close agreement with the findings of Sreekanth *et al.* (2006) and Smita *et al.* (2012), who reported greater plant spread under early planting schedules due to extended vegetative growth periods and optimal climatic conditions.

The maximum number of primary branches (11.97) was observed in treatment G₄P₁, where calendula seedlings were transplanted in the second fortnight of October using the cocopeat-based growing medium (Cocopeat: Sand: FYM: Vermicompost – 2:1:1:1). This enhanced branching may be attributed to the improved physical properties of cocopeat, such as higher porosity and water retention capacity, coupled with the nutrient-rich profile of vermicompost, which collectively enhance nutrient uptake and support vigorous vegetative growth. These observations are in agreement with the findings of Nair *et al.* (2015), who reported that a growing medium composed of cocopeat, sand, FYM, and vermicompost significantly improved the number of branches and flower yield per plant. The enhancement in vegetative and reproductive parameters is likely due to the better availability of essential nutrients particularly nitrogen, phosphorus, and potassium in such organic-rich, soilless media, as previously highlighted by Gupta *et al.* (2007). The superiority of October planting can be attributed to favorable climatic conditions during the early vegetative phase, such as optimal temperature, light, and humidity, which facilitate better cell division and elongation, thereby promoting increased branching. These results suggest that earlier planting extends the vegetative growth period, enabling plants to develop a more robust and well-structured canopy. Similar trends were reported by Laxmi and Pratap (2011) in chrysanthemum and by Hazarika *et al.* (2003), who also observed improved branching and vegetative growth in response to earlier transplanting dates.

Table 2 : Effect of growing media and planting dates on Vegetative growth of Calendula

FACTOR	Plant Height (cm)			Plant spread (cm) E-W			Plant spread (cm) N-S			Number of Primary Branches per plant		
	P1	P2	MEAN	P1	P2	MEAN	P1	P2	MEAN	P1	P2	MEAN
G1	12.08	10.88	11.48	20.71	19.37	20.04	21.36	19.96	20.66	10.38	8.74	9.56
G2	11.96	12.50	12.23	21.23	22.11	21.67	21.62	21.74	21.68	9.21	10.26	9.73
G3	12.37	11.66	12.02	21.44	20.72	21.08	21.65	20.57	21.11	10.00	9.63	9.82

G4	14.44	13.67	14.05	24.45	23.46	23.95	24.41	23.42	23.92	11.97	11.25	11.61
G5	11.61	12.43	12.02	20.82	21.12	20.97	20.76	21.25	21.00	11.19	11.22	11.20
G6	13.01	11.35	12.18	22.59	21.30	21.95	23.01	22.56	22.79	11.69	10.62	11.15
MEAN P	12.58	12.0		21.87	21.35		22.13	21.58		10.7	10.28	
	C.D.	SE(d)	SE(m)	C.D.	SE(d)	SE(m)	C.D.	SE(d)	SE(m)	C.D.	SE(d)	SE(m)
G	0.80	0.38	0.27	0.78	0.37	0.26	0.63	0.30	0.21	0.70	0.33	0.24
P	0.46	0.22	0.15	0.45	0.21	0.15	0.36	0.17	0.12	0.40	0.19	0.13
G X P	1.14	0.55	0.38	1.10	0.53	0.37	0.89	0.43	0.30	0.99	0.48	0.33

As mentioned in Tab 3. The earliest flowering was recorded in treatment G₄P₁, comprising Cocopeat: Sand: FYM: Vermicompost (2:1:1:1) planted in the second fortnight of October, with a value of 62.83 days. The early onset of flowering in this treatment may be attributed to the favorable properties of cocopeat, such as its lightweight structure, excellent aeration, and high water-holding capacity, which collectively enhance root development and overall plant vigor. These results are in agreement with the findings of Treder (2008), who reported earlier flowering in lilies grown in cocopeat-based media. Similar conclusions were drawn by Treder and Nowak (2002), who emphasized the advantages of soilless substrates like cocopeat in promoting early flowering in ornamental crops. The earlier flowering observed under October planting may be due to more favorable environmental conditions, particularly lower ambient temperatures and shorter photoperiods during the early stages of plant development, which can stimulate floral induction in certain long-day species like calendula. These findings are in close conformity with the results reported by Lakshmi *et al.* (2014) in marigold and Sharma *et al.* (2015) in *Gaillardia*, where early planting led to significantly reduced days to flowering, attributed to optimal climatic conditions during the early growth phase. The highest flower count per plant (43.90) was noted in treatment G₄P₁, where seedlings were planted in the second fortnight of October using the cocopeat-based medium. This was closely followed by treatment G₆P₁, comprising Sphagnum Moss: Sand: FYM: Vermicompost (2:1:1:1), which recorded 43.99 flowers per plant. These findings emphasize the superior performance of both cocopeat- and sphagnum-based soilless media when supplemented with FYM and vermicompost, likely due to enhanced nutrient availability and improved aeration in the root zone. The present results are corroborated by Riaz *et al.* (2015), who reported increased flower production with the incorporation of FYM, and by Kameswari *et al.* (2014), who observed that media enriched with cocopeat and FYM significantly improved floral yield. The beneficial effect of these media may be attributed to their capacity to retain moisture, supply essential nutrients, and foster a favorable micro environment for

root and shoot development. The enhanced floral output in October plantings can be attributed to a longer vegetative growth period and a higher number of primary branches, which provided more potential sites for flower initiation. Favorable climatic conditions during early growth stages also likely contributed to better floral differentiation and development. These findings are consistent with those reported by Sreekanth *et al.* (2006) and Ghosh and Pal (2008), who found that earlier planting dates resulted in significantly higher flower production, owing to improved environmental conditions and prolonged growth duration. In Tab. 3 with respect to flower size, indicating that the combined effect of these factors did not influence flower diameter in a statistically meaningful way. However, among all treatment combinations, the largest flower size (5.95 cm) was recorded in G₆P₁, comprising Sphagnum Peatmoss: Sand: FYM: Vermicompost (2:1:1:1) planted in the second fortnight of October. The enhanced flower size observed in this treatment may be attributed to the beneficial physical and biochemical properties of sphagnum peatmoss, including its superior water retention capacity, aeration, and ability to stimulate hormonal activity particularly growth regulators like cytokinins and gibberellins that are crucial for flower and fruit development. These findings align with those of Moneruzzaman *et al.* (2013), who reported that peat-based substrates positively influence the size and quality of flowers through hormonal and physiological enhancement. The improved flower size under October planting may be due to favorable environmental conditions during the early growth stages which allowed for extended vegetative development and better resource accumulation, ultimately contributing to larger floral organs. These results are consistent with the findings of Lakshmi *et al.* (2014) in African marigold and Sharma *et al.* (2015) in *Gaillardia*, where earlier planting dates significantly improved flower size due to optimal temperature and light conditions during critical growth periods. The maximum flowering duration (73.33 days) was recorded in treatment G₄P₁, comprising Cocopeat: Sand: FYM: Vermicompost (2:1:1:1) planted in the second fortnight of October. The superior performance of this treatment

can be attributed to the favorable characteristics of cocopeat, which ensures optimal moisture retention and aeration, and vermicompost, which enriches the medium with essential macro- and micronutrients. This combination creates an ideal root environment that supports sustained flower production over an extended period. These results are in accordance with the findings of Thakur and Grewal (2019) in chrysanthemum cv. 'Kikiobiory', where a growing medium containing cocopeat and FYM significantly extended the flowering period due to improved substrate aeration, moisture retention, and nutrient supply. The prolonged bloom period under October

planting may be attributed to favorable climatic conditions during the early growth phase, which enhanced vegetative vigor and led to the development of robust floral structures capable of sustaining extended reproductive activity. The longer duration of flowering in early-planted calendula is also likely due to greater assimilate accumulation and better resource partitioning toward reproductive organs. These findings are supported by Lakshmi *et al.* (2014) in African marigold, where early transplanting resulted in prolonged blooming owing to improved physiological conditions and extended growth cycles.

Table 3 : Effect of growing media and planting dates on Flowering of Calendula

FACTOR	Days to 50 % Flowering			Number of Flowers			Flower size (cm)			Duration of Flowering		
	P1	P2	MEAN	P1	P2	MEAN	P1	P2	MEAN	P1	P2	MEAN
G1	78.27	81.93	80.09	27.54	26.06	26.80	5.20	4.68	4.94	55.55	51.88	53.72
G2	71.67	75.48	73.57	35.18	33.03	34.12	5.58	5.25	5.41	61.44	59.22	60.33
G3	77.26	73.08	75.17	33.23	35.09	34.16	5.66	5.49	5.57	60.44	56.33	58.38
G4	62.83	67.00	64.91	43.99	41.89	42.94	5.93	5.72	5.83	73.33	68.44	70.88
G5	72.59	72.63	72.61	27.37	28.53	27.97	5.74	5.40	5.57	62.78	67.77	65.27
G6	70.12	72.16	71.14	38.89	35.30	37.09	5.95	5.78	5.86	67.89	62.55	65.22
MEAN P	72.12	73.714		34.37	33.32		5.68	5.38		63.57	61.03	
	C.D.	SE(d)	SE(m)	C.D.	SE(d)	SE(m)	C.D.	SE(d)	SE(m)	C.D.	SE(d)	SE(m)
G	2.67	1.29	0.91	1.22	0.58	0.41	0.40	0.19	0.13	2.64	1.27	0.90
P	1.54	0.74	0.52	0.70	0.34	0.24	0.23	0.11	0.07	1.52	0.73	0.52
G X P	3.78	1.82	1.29	1.72	0.83	0.58	N/A	0.27	0.19	3.74	1.80	1.27

Conclusion

The present investigation clearly demonstrated that both growing media composition and planting dates had a significant impact on the vegetative and flowering performance of calendula. Among the treatments, the growing medium comprising Cocopeat: Sand: FYM: Vermicompost (2:1:1:1) (G₄), particularly when combined with the second fortnight of October planting (P₁), consistently outperformed all other treatments across most parameters. This combination (G₄P₁) resulted in the tallest plants, maximum plant spread, highest number of primary branches, earliest flowering, maximum number of flowers, and longest flowering duration while G₆P₁ produces larger flower size. The superior performance of the treatment (G₄P₁) can be attributed to the enhanced physical, chemical, and biological properties of the medium—such as better aeration, moisture retention, and nutrient availability—as well as the favorable climatic conditions prevailing during early October, which promoted robust vegetative growth and efficient reproductive development. The findings suggest that for optimal growth and floral yield in calendula under Bundelkhand conditions, a growing medium enriched with cocopeat and organic amendments, combined

with early planting in October, offers the most promising results.

References

- Abad, M., Noguera, P. and Bures, S. (2001). National inventory of organic wastes for use as growing media for ornamental potted plant production: case study in Spain. *Bioresour. Technol.*, **77**: 197-200.
- Beckmann-Cavalcante, M.Z., Pivetta, K.F.L., Meinken, E. and Roeber, R. (2009). Growth of *Chrysanthemum × grandiflorum* in different peats and change of peat properties during cultivation. *Acta Horticulturae*, **819**: 181-84.
- Deogade, A.S., Ningot, E.P., Thakare, A.A., Ingole, A.R. and Dahale, M.H. (2020). Effect of potting mixture and pot size on growth and flowering of calendula. *Journal of Pharmacognosy and Phytochemistry*, **9**(6): 1147-51.
- Dubey, R.K., Singh, S., Kukal, S.S. and Kalsi, H.S. (2013). Evaluation of different organic growing media for growth and flowering of petunia. *Communications in Soil Science and Plant Analysis*, **44**: 1777-85.
- Ghosh, P. and Pal, P. (2008). Performance of *Tagetes erecta* L. as influenced by planting time and spacing under West Bengal conditions. *Natural Product Radiance*, **7**(5): 437-43.
- Gupta, Y.C., Dien, L.C., Dhiman, S.R. and Jain, R. (2004). Standardization of growing media under protected environment in gerbera in mid-hills of Himachal Pradesh. *Journal of Ornamental Horticulture*, **7**: 99-102.

- Hazarika, K., Preethi, H., Paswan, L. and Kalitha, P. (2003). Comparison of date of planting and growing conditions on growth and flowering of African marigold (*Tagetes erecta* L.). National Symposium on Recent Advances in Indian Floriculture 13.
- Kameswari, P., Lalitha, G., Girwani, A. and Padmavathamma, A.S. (2014). Effect of different potting media mixtures on growth and flowering of chrysanthemum (*Dendranthema grandiflora* T.). *Progressive Horticulture*, **46**(2): 314-18.
- Lakshmi, R.K.P., Dogra, S., Laishram, N., Bhat, D., Singh, A. and Jamwal, S. (2014). Studies on effect of planting dates and spacing in African marigold (*Tagetes erecta* L.). *Progressive Horticulture*, **46**: 149-52.
- Laxmi, P. and Pratap, M. (2011). Effect of dates of planting on growth, flowering and yield of different Chrysanthemum (*Dendranthema grandiflora* L.) cultivars. *Research on Crops*, **12**(3): 813-16.
- Moneruzzaman, K.M., Boyce, A.N., Normaniza, O., Fruq, G., Rahman, M.M. and Azirun, M.S. (2013). Fruit development, pigmentation and biochemical properties of wax apple as affected by localized application of GA3. *Brazilian Archives of Biology and Technology*, **56**: 11-20.
- Muhammad, A., Muhammad, Q. and Ghulam, M. (2001). Effect of planting dates on growth, flowering and corm characteristics of tuberose (*Polianthes tuberosa*) cv. Single. *International Journal of Agriculture and Biology* **3**: 391-93.
- Nair, S.R., Kumar, K.G. and Santha, K.K. (1985). Effect of planting time and spacing on growth, flower production and seed yield in marigold. *Orissa Journal of Horticulture* **13**: 14-20.
- Nair, S.A. and Bharathi, U. (2015). Influence of potting media composition on pot mum production. *The Bioscan* **10**: 73-76.
- Riaz, A., Arshad, M., Younis, A., Raza, A. and Hameed, M. (2008). Effect of different growing media on the growth and flowering of *Zinnia elegans* cv. Blue Point. *Pak. J. Bot.*, **40**: 1579-1585
- Smita, P., Golliwar, V.J., Panchabhai, D.M., Prarthana, J. and Sonali, D. (2012). Response of African marigold varieties to different planting time on growth and yield under Vidarbha conditions (Maharashtra). *Journal of Soils and Crops* **22**(1): 183-87.
- Sreekanth, P., Padma, M., Chandrasekhar, R. and Madhulety, T.Y. (2006). Effect of planting time, spacing and nitrogen levels on flowering of African marigold (*Tagetes erecta* L.). *Journal of Ornamental Horticulture*, **9**(2): 97-101.
- Thakur, T. and Grewal, H.S. (2019). Influence of potting media compositions on flower production of chrysanthemum (*Chrysanthemum morifolium* Ramat) cultivar 'Kikiobiory'. *Journal of Plant Nutrition*, **42**: 1861-67.
- Thumar, V.B., Chovatiya, V.N., Butani, M.A., Bhalu, B.V. and Chavda, R.J. (2020). Effect of potting media on growth and flower yield of rose (*Rosa hybrida* L.) cv. Top Secret under protected condition. *Journal of Pharmacognosy and Phytochemistry* **9**(2): 408-10.
- Treder, J. and Nowak, J. (2002). The use of coconut substrate in the cultivation of bedding plants. *Journal of Agricultural Science*, **485**: 335-58.
- Treder, J. (2008). The effects of cocopeat and fertilization on the growth and flowering of oriental lily 'Star Gazer'. *Journal of Fruit and Ornamental Plant Research*, **16**: 361-70.
- Vendrame, A.W., Maguire, I. and Moore, K.K. (2005). Growth of selected bedding plants as affected by different compost percentages. *Proceedings of the Florida State Horticultural Society*, **118**: 368-71.
- Verma, K. (2001). Effect of planting dates, pinching methods and foliar application of nitrogen on growth and flowering of carnation under protected conditions. M.Sc. Thesis submitted to Dr. Y.S. Parmar University of Horticulture and Forestry, Nauni, Solan.