



# Plant Archives

Journal homepage: <http://www.plantarchives.org>

DOI Url : <https://doi.org/10.51470/PLANTARCHIVES.2026.v26.no.1.213>

## IDENTIFICATION, PURIFICATION AND CHARACTERIZATION OF COMPOUND ISOLATED FROM *PSORALIA CORYLIFOLIA* USING A NOVEL APPROACH

Shri Ram Bharat<sup>1</sup>, Suman Kumar Jha<sup>2</sup>, Rajeev Saini<sup>3</sup>, Satyavrat Fort<sup>1</sup>, Brijesh Kumar<sup>1</sup>,  
Rishabh Kumar<sup>3</sup> and Archana Suyal<sup>3\*</sup>

<sup>1</sup>Patanjali Ayurveda Limited, Padartha, Haridwar, Uttarakhand, India.

<sup>2</sup>Patanjali Foods Limited, Padartha, Haridwar, Uttarakhand, India.

<sup>3</sup>R & D, Patanjali Natural Coloroma Private Limited, Padartha, Haridwar, Uttarakhand, India.

\*Corresponding author E-mail : [archana.suyal@patanjalicoloroma.com](mailto:archana.suyal@patanjalicoloroma.com)

(Date of Receiving-01-12-2025; Date of Revision-18-01-2026; Date of Acceptance-17-02-2026)

### ABSTRACT

*Psoralia corylifolia* Linn. (Leguminosae) is one of the important pharmaceutical potent plants possesses various activities like antibacterial, anticancer, antioxidant, antifungal, anti-depressant and antitumor, therefore used in different medicinal formulations worldwide. This plant has given important lead in drug research due to its scientific finding, resulting in the discovery of various novel molecules. The objective of the present study was to develop a flash chromatography method for isolation and purification of psoralen biomolecule from *Psoralia corylifolia* Linn. seeds extract and its further identification and characterization using different spectral analytical techniques. Absorption maxima of psoralen at 247 nm was confirmed using UV spectrophotometer. TLC and HPLC quantification of psoralea were done before and after isolation from seeds extract of *Psoralia corylifolia* Linn. Flash chromatography method were used for purification and isolation. About 250 g of raw material yielded 15.63 g of crude hexane extract. Further using flash chromatography about 68 mg of white needle-shaped crystalline powder of psoralea of over 98% purity, determined by high performance liquid chromatography (HPLC) was obtained. The identification and characterization of psoralea performed using FTIR, <sup>1</sup>H NMR and <sup>13</sup>C NMR spectral data.

**Key words :** *Psoralia corylifolia*, Flash Chromatography, High Performance Liquid Chromatography (HPLC).

### Introduction

*Psoralia corylifolia* Linn. is a traditional medicinal herb, belongs to Leguminosae family, commonly known as “Bakuchi” and ‘Bu GhuZhi’ in Indian traditional medicine and in traditional Chinese medicine respectively (Patel, 2023; Alam *et al.*, 2018). *Psoralea corylifolia* Linn., has diversified uses in agriculture, ayurvedic as well as allopathic system of medicines with thousands of years of clinical application. Most of the therapeutically active components are present in its seeds or fruits are coumarins (psoralidin, psoralen, isopsoralidin, isopsoralen, angelicin and corylifolin), flavonoids namely neobavaisoflavone, isobavachalcone, bavachalcone, bavachinin, bavachin, corylin, corylifol, corylifolin and 6-prenylnaringenin and meroterpenes (3-hydroxybakuchiol

and bakuchiol). Its essential oil is rich in limonene,  $\alpha$ -elemene,  $\gamma$ -elemene,  $\beta$ -caryophyllenoxide, 4-terpineol, linalool, geranylacetate (Koul *et al.*, 2019; Bertoli *et al.*, 2004). *Psoralea corylifolia* found throughout India in Himalayas, Uttarakhand, adjoining area of Uttar Pradesh, semi-arid regions of Punjab and Rajasthan, Bundelkhand, Bengal and Bombay, Deccan and Karnataka (Vidua *et al.*, 2015). The extracts and active components of *Psoralea corylifolia* Linn. demonstrated multiple biological activities, including anticancer, estrogenic, antitumor, anti-oxidant, anti-microbial, anti-depressant, anti-inflammatory and osteoblastic and hepatoprotective activities (Zhang *et al.*, 2016; Pandey *et al.*, 2013). The principal bioactive compound of bakuchi is psoralen, which is a tricyclic coumarin like aromatic compound, having

molecular structure 7H-Furo [3, 2-g] benzopyran-7-one (Pathak *et al.*, 1959). *Psoralea corylifolia* Linn. biomolecule psoralen (coumarins) widely used as quality control components in various herbal formulations. Psoralen which on exposure to sun forms melanin in depigmented skin, found to intercalate into DNA covalently, where on exposure to long wave UV light form mono- and di- adducts and thus are used for the treatment of leucoderma a hypo pigmented lesions of the skin and vitiligo an autoimmune disease (Sui *et al.*, 2020). The isolation and purification of *Psoralea* and related compounds from *Psoralea corylifolia* Linn., reported methods are tedious and usually require highly sophisticated instruments which are not cost effective. Due to the important biological properties and broad applications of psoralen, we have attempted to develop an efficient method to isolate and purify the pure compound from the extract of *Psoralea corylifolia* Linn. seeds. The subsequent preparative isolation of compounds of interest used here is flash chromatography (flash purification). It enables rapid separation, reliable and expeditious approach for isolation of pure compounds from complex natural product mixtures, as it is hybrid of gravity column chromatography and preparative liquid chromatography, in this solvent coupled with a pressurized gas like nitrogen in order to assist in the separation process (Roge *et al.*, 2011; Girijal *et al.*, 2018). The advantage of flash chromatography method over other crude purification techniques is that it has ability to separate broad variety of pure compounds more efficiently, offering rapid and finest results (Kustrin *et al.*, 2022). This study focus on flash chromatography method development to separate and isolate high purity bioactive compound psoralen from *Psoralea corylifolia* Linn. extract in the most cost-efficient way and its further characterization via FTIR, NMR spectroscopy, guided by thin-layer chromatography (TLC) and high performance liquid chromatography (HPLC). In addition, this study gives a comprehensive overview of psoralen pharmacological potential, which will be helpful in further research.

### Materials and Methods

**Bakuchi plant seeds :** The seeds of *Psoralea corylifolia* Linn. were collected from local vendor, Haridwar. Authenticity of raw seeds were confirmed by Patanjali Research Foundation Herbarium (PRHF), Haridwar. The seeds were air dried in shadow, reduced to coarse powder with grinder and stored in airtight zipper bag till further use.

**Chemicals :** All solvent used were of reagent grade and analytical grade purchased from Merck Ltd, Mumbai,

India. Reference standards of psoralen (purity  $\geq 99.0\%$ ) was purchased from Sigma Aldrich, Bangalore, India.

**Extraction of crude *Psoralea* :** Air dried powder (250 g) of *Psoralea corylifolia* Linn. seeds was extracted successively by soxhletion (3 times, each for 3 hours,  $50 \pm 5^\circ\text{C}$ ), using n-hexane. The extraction solvents used were removed under vacuum pressure using rotary evaporator. The extracts were collected weighed 15.62 g and their yield percentage were recorded and was thereafter, stored under cool temperature until further analysed.

### Instrumentation

#### Determination of absorption maxima

**UV spectrophotometer :** The measurement of the UV spectra and absorbance readings was conducted using the Shimadzu UV- 1800 model, UV-Visible Double Beamed Spectrophotometer with bandwidth 1 nm, wavelength accuracy 0.1 nm and quartz cuvettes with 1 cm path length, over the specific  $\lambda_{\text{max}}$  at 200-400 nm. The glassware used during the process were treated with a chromic acid and sulphuric acid solution, further rinsed thoroughly with double distilled water and dried in the hot air oven. Accurately weighed 10 mg of psoralen, transferred to 100 ml volumetric flask. Volume makeup to 100 ml by adding methanol in order to get the concentration of  $100\mu\text{g/ml}$ . The resultant solution ( $100\mu\text{g/ml}$ ) were scanned on UV spectrophotometer so as to observe the absorption maxima of psoralen.

#### Identification and Quantification

**TLC (Thin layer chromatography) :** Method for qualitative determination of psoralen in solvent extracted *Psoralea corylifolia* Linn. seeds extract and isolated compound were determined. The advantage of using thin-layer chromatography (TLC) method as large number of samples can be simultaneously analyzed using small volume of mobile phase thus lowering analysis time and cost per analysis. Precoated (Merck) analytical plates with fluorescent indicator (silica gel 60 F<sub>254</sub>, aluminum sheets) were used for thin layer chromatography (TLC). Glass capillary tubes were used for spotting in duplicate. One mg psoralen RS in 10 ml volumetric flask. Add 5 ml methanol dissolve and makeup the solution. For sample preparation 1 gm crude extract was dissolved in methanol in 25 ml volumetric flask. The plate was developed in developing chamber, pre-saturated for 2 hours with mobile phase hexane: ethyl acetate, (7:3 v/v) (solvent front 6.0 cm). The plates were observed at wavelength 254 nm and 366 nm. Retention factor ( $R_f$  value) for reference standard and sample were calculated.

### HPLC (High performance liquid chromatography)

Psoralea was simultaneously determined in *Psoralea corylifolia* Linn. seeds extract using HPLC system (Shimadzu *i*-series LC 2030), equipped with a quaternary pump, a variable wavelength UV-VIS detector, RP-18 column (250 mm × 4.6 mm i.d., particle size 5 $\mu$ ). The mobile phase ratio A: Mixture of 0.146 g KH<sub>2</sub>PO<sub>4</sub> / 0.5 ml OPA in 1000 ml milli Q water, mobile phase B: 100 % Acetonitrile (v/v). Gradient phase used is shown in Fig. 1. The flow rate was 1.5 ml/ min, psoralea was detected at wavelength of 247 nm at room temperature 25°C. The retention time of psoralea reference standard was 13.397 min under the above conditions. Data were collected and processed using computing integrator chromatography workstation (Labsolution software).

**Standard preparation :** 100 mg reference standard of psoralen was dissolved in 100ml methanol, sonicated for 15min. Further 10ml was taken and diluted with 100ml methanol.

**Sample preparation :** One gm extract sample was dissolved in methanol in 100ml volumetric flask. Further 10ml was taken and diluted with 100ml methanol and filtered through 0.45 $\mu$ m millipore filter prior to HPLC analysis.

### Flash Chromatography

**Preparative isolation and purification :** *Psoralea corylifolia* Linn. seeds extract containing psoralea was purified using Flash (Buchi, C- 850 Flash Prep) shown in Fig. 3, integrated with a variable wavelength UV-VIS detector and ELSD, Pressure range flash mode: 0-50 bar and prep mode: 0-300 bar. Flash cartridges (60-120

mesh) column. The solvent used were hexane: ethyl acetate (v/v) in gradient mode depicted in Fig. 2. Flow rate was at 25 mL/min. Variable wavelength were used in range  $\lambda_1$  - 210,  $\lambda_2$  - 247,  $\lambda_3$  - 310,  $\lambda_4$  - 420 at room temperature 25°C.

**Sample preparation for loading :** Ten gm hexane extracted *Psoralea corylifolia* seeds extract was added into 30gm silica gel (60-120 mesh size), mixed properly by gradually adding small amount of hexane. Then dried in oven at 50-60°C till solvent evaporates completely. Further packed under vacuum in flash empty solid loader.

### Characterization

**FTIR (Frontier Transform Infrared spectroscopy) :** The characterization of isolated compound was carried out using FTIR (Shimadzu IR). The testing was carried out by using KBr pellet method. KBr was dried in the oven at 40°C and it was then mixed with the sample. The sample was compressed using a pellet press. The sample was placed in IR sample holder. A background scan was taken followed by the IR scan of the sample in range of 500- 4000 cm<sup>-1</sup>

**NMR (Nuclear magnetic resonance, <sup>1</sup>H and <sup>13</sup>C) :** <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra of preparative isolated compound were measured at 500 MHz and 125 MHz respectively on a Bruker avance NEO 500 spectrophotometer, CDCl<sub>3</sub> with or without tetramethylsilane as an internal standard. Chemical shifts are given in  $\delta$  (ppm).

### Results and Discussion

The identification and characterization of the compounds usually involves a combination of different spectral optimized techniques including UV, FTIR, <sup>1</sup>H and <sup>13</sup>C NMR. Under the TLC chromatographic conditions, the retention factor ( $R_f$  value) for reference standard, crude extract and isolated compound were found to be  $0.72 \pm 0.02$ , which indicates the presence of psoralen. The characteristic retention time ( $R_t$ ) chromatograms of the psoralen standard and crude extract of seed of *Psoralea corylifolia* Linn. are shown in Figs. 3 and 4, respectively, in which the  $R_t$  of psoralen reference standard was obtained at 14.397 and  $R_t$  of psoralen in crude extract of *Psoralea corylifolia* Linn. seeds were obtained at 14.395. Further, shown in Fig. 4, quantitative HPLC analysis of the crude extract from *Psoralea corylifolia* Linn.




Fig. 1 : Gradient mobile phase method for Hplc.




Fig. 2 : Gradient mobile phase method for flash chromatography.




Fig. 3 : Chromatogram of psoralen reference standard.




Fig. 4 : Chromatogram of *Psoralea corylifolia* seeds crude extract.

seeds contained psoralen a major compound, purity based on HPLC peak area percentage in crude extract was reported 49.3%.

Until now, no studies have been reported of psoralen isolation by flash chromatography. Past literature data has reported isolation of psoralen by column chromatography (Rajput *et al.*, 2008; Lu *et al.*, 2014). Crude *Psoralea corylifolia* extract containing psoralen was loaded on flash chromatography and first eluent hexane (100%) and ethyl acetate (0%) v/v after 30 min elute fraction consisted of yellow brown colour liquid characterized as Bakuchiol (50 mg), while the fraction eluted with hexane and ethyl acetate (95:05 v/v) at 1 hour 27 min consisted of white needle-shaped crystalline powder (68 gm) which was characterized as psoralen (>98%) by comparing with reference standard using TLC and HPLC, shown in Fig. 5.

The isolated psoralen solution 100 µg/ml were scanned on UV spectrophotometer so as to observe the absorption maxima. The UV spectrum exhibited absorption maxima at  $\lambda_{max}$  245 nm and 300 nm, consistent with the psoralen chromophore, shown in Fig. 6. Further, structural identification and characterization of purified psoralen carried out by FTIR,  $^1\text{H}$  NMR and

$^{13}\text{C}$  NMR.

#### Identification and Characterization of isolated compound using FTIR

Fourier Transform Infrared Spectroscopy (FTIR) serves as a fingerprint technique, helps characterize the functional groups present in isolated compound, depicted absorption bands in Fig. 7. The unique pattern of peaks allow identification and confirm presence of psoralen (Rajput *et al.*, 2008). FTIR analysis is important technique for understanding the chemical properties and potential medicinal uses of psoralen.

**Aromatic C-H Stretch:** Peaks at approximately 3155 and 3060  $\text{cm}^{-1}$  indicate the presence of aromatic C-H bonds. **C=O Stretch (Lactone):** A strong peak around 1712  $\text{cm}^{-1}$  signifies the carbonyl (C=O) group in unsaturated lactone ring. **C=C Ring Stretch:** Absorption at approximately 1634  $\text{cm}^{-1}$  is attributed to the C=C bonds within the fused rings. **C-O Stretch:** Absorption around 1131  $\text{cm}^{-1}$  is characteristic of C-O stretching vibrations. **C-O-C Ether Group:** The peak at about 1092  $\text{cm}^{-1}$  corresponds to the C-O-C ether linkage. **Out-of-plane C-H Bend:** A peak at approximately 896, 823 and 717  $\text{cm}^{-1}$  is associated with the bending of the C-H bonds. The above data clearly indicates that the isolated




Fig. 5 : Chromatogram of flash isolated psoralen.




Fig. 6 : Chromatogram of psoralen UV spectrum.




Fig. 7 : FTIR of isolated compound.




Fig. 8 : <sup>1</sup>H NMR of isolated compound.

compound is psoralen as it contains carbonyl groups and lactone bonding arrangements in a ring fused to a coumarin.

**Identification and characterization of isolated compound by <sup>1</sup>H NMR and <sup>13</sup>C NMR :** The flash isolated white needles crystal with MeOH; mp 162-166°C was characterized by NMR spectroscopy. The <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) spectrum showed characteristic signals (500 MHz, CDC13) δ, 6.36 (<sup>1</sup>H, d, J= 9.6 Hz, H-3), 6.81 (<sup>1</sup>H, d, J = 2.4 Hz, H-11), 7.46 (<sup>1</sup>H, s, H-8), 7.66 (<sup>1</sup>H, s, H-5), 7.67 (<sup>1</sup>H, d, J = 2.4 Hz, H-12), 7.79 (<sup>1</sup>H, d, J = 9.6 Hz, H-4); EIMS m/z (relative intensity) 186 (100, [M]<sup>+</sup>), 158, 130, 102.

The psoralen <sup>13</sup>C-NMR, (100.5MHz, CDC13) spectrum exhibited characteristic signals at δ (ppm): 161.0 (2-C), 114.6 (3-C), 144.0 (4-C), 119.8 (5-C), 124.8 (6-C), 156.4 (7-C), 99.8 (8-C), 152.0 (9-C), 115.4 (10-C), 146.9 (2'-C), 106.3 (3'-C), confirming the psoralen structure.

### Conclusion

The optimized flash chromatography was efficiently used for the preparative isolation and purification of marker compound, psoralen from *Psoralea corylifolia* Linn. seeds. High-purity (>98%) psoralen was obtained from crude extract in one step separation that can be used for further scientific studies. The separation and yields of pure psoralen obtained by this method are reproducible. Further, identified and comprehensive characterized as psoralen via FTIR and NMR spectral data was done. The separation and yields of pure psoralen obtained by this method are reproducible. This streamlined flash chromatography approach enables isolation of biomolecule psoralen which have remarkable medicinal benefits include anti-inflammatory, antimicrobial and anticancer properties. Therefore, invariably on high demand in pharmaceutical, cosmetic, herbal and other related industries.

### Acknowledgements

The authors would like to acknowledge Patanjali Ayurved Limited and Patanjali Foods Limited for providing all necessary support in preparation of this manuscript.

### References

Alam, F., Khan G.N. and Asad M.H.H.B. (2018). *Psoralea corylifolia* L: Ethnobotanical, biological and chemical aspects: A review. *Phytother Res.*, **32**(4), 597-615. doi: 10.1002/ptr.6006. Epub 2017 Dec 15. PMID: 29243333; PMCID: PMC7167735.

Bertoli, A., Menichini F., Noccioli C., Morelli I. and Pistelli L. (2004). Volatile constituents of different organs of

*Psoralea bituminosa* L. *Flav Frag J.*, **19**, 166-171. doi: 10.1002/ffj.1315.

Girija, D., Padma A. and Reddy B.V. (2018). A Review on Flash Chromatography. *Int. J. Chem. Pharm. Sci.*, **6**(11), 300-305. [https://www.researchgate.net/publication/381963650\\_A\\_Review\\_on\\_Flash\\_Chromatography](https://www.researchgate.net/publication/381963650_A_Review_on_Flash_Chromatography).

Koul, B., Taak P., Kumar A., Kumar A. and Sanyal I. (2019). Genus *Psoralea*: A review of the traditional and modern uses, phytochemistry and pharmacology. *J Ethnopharmacol.*, **25**; **232**, 201-226. doi: 10.1016/j.jep.2018.11.036. Epub 2018 Dec 3. PMID: 30521980; PMCID: PMC7127090.

Kustrin, S.A. Gegechkori V., Mohammed E.U.R., Ku H. and Morton D.W. (2022). Isolation of Bioactive Pentacyclic Triterpenoid acids from Olive Tree Leaves with Flash Chromatography. *Appl. Sci.*, **12**(3), 996. <https://doi.org/10.3390/app12030996>.

Lu, H., Zhang L., Liu D., Tang P. and Song F. (2014). Isolation and purification of psoralen and isopsoralen and their efficacy and safety in the treatment of osteosarcoma in nude rats. *Afr. Health Sci.*, **14**(3), 641-7. doi: 10.4314/ahs.v14i3.20. PMID: 25352883; PMCID: PMC4209630.

Pandey, P., Mehta R. and Upadhyay R. (2013). Physico-chemical and preliminary phytochemical screening of *Psoralea corylifolia*. *Arch Appl. Sci. Res.*, **5**, 261-265.

Patel, D.K. (2023). An Overview of Biological Importance, Pharmacological Aspects, and Analytical Techniques of Bavachin from *Psoralea corylifolia* Linn. *Current Functional Foods*, **1**(1), 55-63. [10.2174/2666862901666220411123932](https://doi.org/10.2174/2666862901666220411123932)

Pathak, M.A. and Fitzpatrick T.B. (1959). The nature of the psoralen- DNA photoaddition reaction. *J. Investigative Dermatol.*, **32**(2), 255-262. doi: 10.1038/jid.1959.38

Rajput, S.J., Zade V. and Rai P. (2008). Studies on extraction, isolation and estimation of psoralen from the fruits of *Psoralea corylifolia*. *Phcog. Mag.*, **4**, 52-56. <https://phcog.com/article/assets/v4/i13/PhcogMag-4-13-52.pdf>.

Roge, A.B., Firke S.N., Kawade R.M., Sarje S.R. and Vadvalkar S.M. (2011). Brief review on: flash chromatography. *IJPSR*, **2**(8), 1930-1937. [http://dx.doi.org/10.13040/IJPSR.0975-8232.2\(8\).1930-37](http://dx.doi.org/10.13040/IJPSR.0975-8232.2(8).1930-37).

Sui, X., Liu T., Liu J., Zhang J., Zhang H. and Wang H. (2020). Ultrasonic-enhanced surface-active ionic liquid-based extraction and defoaming for the extraction of psoralen and isopsoralen from *Psoralea corylifolia* seeds. *Ultrason. Sonochem.*, **69**, 105263. doi: 10.1016/j.ultsonch.2020.105263.

Vidua, G., Shrivastava D., Tiwari A.K., Bhardwaj S., Tripathi M. and Tiwari A. (2015). Quality control and pharmacognostical activity of Linn. *Psoralea corylifolia*. *J. Chem. Pharm. Res.*, **7**(6), 249-256.

Zhang, X., Zhao W., Wang Y., Lu J. and Chen X. (2016). The Chemical Constituents and Bioactivities of *Psoralea corylifolia* Linn.: A Review. *Am J. Chin. Med.*, **44**(1), 35-60. doi: 10.1142/S0192415X16500038. PMID: 26916913.