Plant Archives Vol. 26, No. 2, 2026 pp. 2245-2250 e-ISSN:2581-6063 (online), ISSN:0972-5210

Plant Archives

Journal homepage: http://www.plantarchives.org
DOI Url : https://doi.org/10.51470/PLANTARCHIVES.2026.v26.no.2.308

POLLINATORS UNDER PRESSURE: CHALLENGES, STRATEGIES AND FUTURE DIRECTIONS: A REVIEW

P. Srikanth[1]*, G. Tirumala Geethika[1], M. Poojith Kumar Reddy[2], T. Thamizharasu[1], S. Nivya[1], R. Likith Kumar[1], Nishrita Khangia[1] and and K. Ganesh[1]

[1]Department of Sericulture, College of Temperate Sericulture, Sher-e-Kashmir University of Agricultural Sciences and Technology, Shalimar, Srinagar, Jammu and Kashmir, India.
[2]Department of Plant Protection - Entomology, College of Horticulture, Anantharajupeta, Railway Koduru, India.
*Corresponding author E-mail : srikanthperumal04@gmail.com
(Date of Receiving-18-05-2026; Date of Revision-11-08-2026; Date of Acceptance-21-08-2026)

ABSTRACT

Pollinators sustain the reproduction of the majority of flowering plants and underpin the productivity of more than three-quarters of global food crops, yet their populations are declining at an alarming rate across continents. Habitat loss and fragmentation, pesticide exposure to neonicotinoids, pathogens and parasites such as *Varroa destructor* and *Nosema ceranae*, climate-driven phenological mismatches, invasive species, and agricultural intensification act both individually and synergistically to reduce pollinator abundance, diversity, and colony persistence. These pressures translate into measurable declines in crop yield, fruit quality, and wild plant reproduction, with cascading effects on ecosystem stability and food security. Habitat restoration, pollinator-friendly farming practices, targeted policy frameworks, disease-resistant breeding programs and technology-assisted monitoring each address specific components of this decline, and evidence indicates that their effectiveness increases substantially when deployed as coordinated, landscape-level interventions rather than isolated measures.

Key words : Climate change, Colony collapse disorder, Ecosystem services, Habitat restoration, Neonicotinoids, Pollinator decline.

Introduction

Pollinators are indispensable components of terrestrial ecosystems, playing a critical role in the reproduction of flowering plants and the maintenance of biodiversity. Globally, it is estimated that over 75% of major food crops depend to some extent on animal-mediated pollination (Klein *et al*., 2007). This ecological service contributes not only to agricultural productivity but also to the stability of natural ecosystems. The economic value of pollination has been estimated at hundreds of billions of dollars annually, underscoring its importance to food security and livelihoods worldwide (Gallai *et al*., 2009). Despite this significance, pollinator populations are experiencing alarming declines, raising concerns about the sustainability of agroecosystems and the resilience of global food systems.

Historical context and importance

The relationship between humans and pollinators dates back thousands of years. Honeybees (*Apis mellifera*) have been domesticated for honey production and crop pollination since antiquity, while wild pollinators such as bumblebees (*Bombus* spp.), solitary bees (*Osmia* spp.), butterflies, moths, beetles, flies, birds and bats have long supported natural and agricultural ecosystems (Michener, 2007). Pollinators enhance crop yields, improve fruit quality, and increase genetic diversity through cross-pollination. Their role extends beyond agriculture, as they sustain wild plant populations that provide habitat and food for other organisms, thereby maintaining ecological balance (Ollerton *et al*., 2011).

Global Decline of Pollinators

In recent decades, evidence has accumulated of

widespread pollinator declines across continents. Studies in Europe and North America have documented reductions in both managed honeybee colonies and wild pollinator species (Biesmeijer *et al*., 2006). Factors contributing to these declines include habitat loss due to agricultural intensification, pesticide exposure, climate change, invasive species, and emerging pathogens (Goulson, 2013). The phenomenon of colony collapse disorder in honeybees, first reported in the mid-2000s, highlighted the vulnerability of pollinators to multiple stressors acting synergistically (van Engelsdorp *et al*., 2009). Similar declines have been observed in bumblebee populations, with several species now listed as threatened or endangered (Cameron *et al*., 2011). The decline of pollinators has profound implications for agriculture and biodiversity. Crops such as apples, almonds, cucurbits, coffee and cocoa are highly dependent on pollinators, and reductions in pollinator abundance can lead to yield losses and decreased quality (Aizen and Harder, 2009). Beyond agriculture, pollinator declines threaten wild plant reproduction, potentially leading to cascading effects on ecosystems and the species that depend on them (Memmott *et al*., 2004). The economic consequences are equally severe, with estimates suggesting that pollinator loss could jeopardize global food production and increase reliance on costly artificial pollination methods.

Drivers of Decline

Pollinator declines are driven by a complex interplay of factors. Habitat loss and fragmentation reduce the availability of floral resources and nesting sites (Kremen *et al*., 2007). Pesticides, particularly neonicotinoids, impair pollinator navigation, foraging, and reproduction (Goulson, 2013). Climate change alters flowering phenology and pollinator activity, leading to mismatches between plants and their pollinators (Aizen and Harder, 2009). Pathogens such as *Nosema ceranae* and parasites like the *Varroa* mite further weaken pollinator populations (Evans and Schwarz, 2011). Invasive species, including non-native pollinators and plants, disrupt native pollination networks (Stout and Morales, 2009).

Need for conservation

Given the ecological and economic importance of pollinators, their conservation has become a global priority. International initiatives such as the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services have emphasized the urgency of protecting pollinators to safeguard food security and biodiversity (IPBES, 2016). Conservation strategies must integrate ecological principles with modern technologies, combining habitat restoration, sustainable farming practices and policy frameworks. Community-based approaches, involving farmers and local stakeholders, are essential to ensure the success of conservation efforts (Kremen and Merenlender, 2018).

Aim and Scope of this Review

This review aims to consolidate current evidence on the principal drivers of global pollinator decline and to critically evaluate the conservation strategies proposed to address them. It draws together ecological, agronomic, technological and policy perspectives to offer an integrated framework for researchers, policymakers and practitioners engaged in pollinator conservation, and to identify priority directions for future research.




Fig. 1 : Overview of pollinator ecosystem services and their dependence on major stressors.

Threats to Pollinators

Pollinators worldwide are facing a convergence of pressures that threaten their survival and the ecosystem services they provide. These threats are not isolated; they often act synergistically, creating cascading effects that accelerate declines (Potts *et al*., 2010).

Habitat Loss and Fragmentation

The expansion of agriculture and urbanization has drastically reduced natural habitats. Conversion of diverse landscapes into monocultures eliminates floral resources and nesting sites, while fragmentation isolates populations and limits gene flow. Studies in Europe demonstrated that wild bee diversity declines sharply in intensively farmed landscapes (Biesmeijer *et al*., 2006). Similarly, Steffan-Dewenter and Tscharntke (1999) showed that habitat isolation reduces pollinator abundance and increases extinction risk.

Pesticide Exposure

Chemical pesticides, particularly neonicotinoids are among the most significant threats. These systemic

insecticides contaminate nectar and pollen, impairing navigation and memory in bees (Goulson, 2013). Whitehorn *et al.* (2012) found that chronic exposure reduced bumblebee colony growth and queen production. Even fungicides and herbicides, though not directly lethal, can interact with insecticides to amplify toxicity (Pettis *et al*., 2013). The reliance on chemical inputs in modern agriculture thus creates a hostile environment for pollinators.

Pathogens and Parasites

Pollinators are increasingly burdened by pathogens and parasites. Honeybees are weakened by *Varroa destructor* mites, which feed on hemolymph and transmit viruses (Rosenkranz *et al*., 2010). Microsporidian parasites such as *Nosema ceranae* reduce bee longevity and foraging efficiency (Evans and Schwarz, 2011). Viral infections, including deformed wing virus, further compromise colony health (Genersch, 2010). Spillover of pathogens from managed bees to wild species has also been documented, raising concerns about disease transmission across pollinator communities (Graystock *et al*., 2013).

Climate change

Climate change alters pollinator distribution, abundance and activity. Rising temperatures and shifting precipitation patterns affect flowering phenology, leading to mismatches between plants and pollinators (Aizen and Harder, 2009). Extreme weather events such as heatwaves and storms reduce survival and disrupt foraging. Kerr *et al*. (2015) reported that bumblebee ranges are contracting, with species failing to expand southward despite warming conditions. These changes destabilize pollination networks and threaten crop yields.

Invasive species

Invasive species disrupt native pollinator communities by competing for floral resources or introducing novel pathogens. Non-native honeybees can outcompete wild bees for nectar and pollen, reducing their reproductive success (Stout and Morales, 2009). Invasive plants alter floral landscapes, sometimes providing poor-quality resources that fail to sustain pollinator populations (Morales and Traveset, 2009). The Asian hornet (*Vespa velutina*), an invasive predator, has caused significant mortality in European honeybee colonies (Monceau *et al*., 2014).

Agricultural Intensification and Monocultures

Modern agriculture often relies on monocultures, which provide abundant but temporally limited floral resources. Once crops finish flowering, pollinators face resource scarcity. Rundlof *et al.* (2015) showed that seed coatings with neonicotinoids negatively affected wild bee populations in monoculture landscapes. Intensive farming also reduces landscape heterogeneity, eliminating hedgerows and wildflower strips that support pollinators. While managed honeybees are often introduced to compensate, this practice increases vulnerability to disease outbreaks and pesticide exposure (Aizen and Harder, 2009).

Synergistic effects

The most concerning aspect of these threats is their tendency to act synergistically. Pesticide exposure can weaken immune systems, making pollinators more susceptible to pathogens (Pettis *et al*., 2013). Habitat loss amplifies the impacts of climate change by reducing refugia. Potts *et al*. (2010) emphasized that multiple stressors create a compounding pressure that accelerates pollinator declines.

Conservation Strategies

Conserving pollinators requires integrated approaches that address the multiple stressors driving their decline. Because threats such as habitat loss, pesticide exposure, pathogens and climate change often act together, strategies must be holistic and adaptable to local contexts (Potts *et al*., 2010).

Habitat Restoration

Restoring and enhancing habitats is one of the most effective strategies. Establishing wildflower strips, hedgerows and buffer zones provides continuous floral resources and nesting sites (Kremen *et al*., 2007). In Europe, agri-environment schemes have successfully increased pollinator abundance by incentivizing farmers to plant diverse vegetation (Carvell *et al*., 2007). Urban initiatives, such as rooftop gardens and pollinator parks, also contribute to habitat restoration, particularly in regions where agricultural land is scarce (Baldock *et al*., 2015).




Fig. 2 : Convergence of major stressors driving global pollinator decline.

Table 1 : Major Threats to Pollinators and their Documented impacts.

Threat	Mechanism	References
Habitat loss and fragmentation	Reduces floral resources, isolates populations, limits gene flow	Biesmeijer *et al.* (2006), Steffan-Dewenter and Tscharntke (1999)
Pesticide exposure (neonicotinoids)	Impairs navigation, memory, colony growth and queen production	Goulson (2013), Whitehorn *et al.* (2012), Pettis *et al.* (2013)
Pathogens and parasites	*Varroa destructor*, *Nosema ceranae* and deformed wing virus weaken colonies	Rosenkranz *et al.* (2010), Evans and Schwarz (2011), Genersch (2010), Graystock *et al.* (2013)
Climate change	Alters flowering phenology, causes plant-pollinator mismatches, contracts species ranges	Aizen and Harder (2009), Kerr *et al*. (2015)
Invasive species	Competes for floral resources, introduces pathogens, predation (e.g., *Vespa velutina*)	Stout and Morales (2009), Morales and Traveset (2009), Monceau *et al.* (2014)
Agricultural intensification and monocultures	Temporally limited floral resources, landscape homogenization	Rundlof *et al.* (2015), Aizen and Harder (2009)

Pollinator-friendly Farming

Agricultural practices can be modified to support pollinators. Reduced tillage preserves soil-nesting bee habitats, while crop rotation and intercropping increase floral diversity (Garibaldi *et al*., 2014). Organic farming systems, which avoid synthetic pesticides, often harbor greater pollinator diversity compared to conventional systems (Hole *et al*., 2005). Integrating Integrated Pest Management principles reduces pesticide use and minimizes harm to beneficial insects (Goulson, 2013).

Policy Frameworks

International and national policies play a crucial role. The Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services emphasized the urgency of coordinated global action (IPBES, 2016). In the European Union, the Pollinator Initiative promotes habitat restoration and pesticide regulation. Similarly, the U.S. Pollinator Partnership works with farmers and communities to implement pollinator-friendly practices. Policies that incentivize conservation through subsidies or certification schemes can encourage widespread adoption (Klein *et al*., 2007).

Genetic approaches

Advances in genetics offer new opportunities. Breeding disease-resistant honeybee strains can reduce vulnerability to pathogens (Evans and Schwarz, 2011). Genomic studies provide insights into pollinator resilience and adaptation to environmental stressors (Zayed, 2009). Conservation of genetic diversity in wild pollinators is equally important, as it enhances population stability and adaptability (Cameron *et al*., 2011).

Technological Monitoring

Modern technologies enhance monitoring and management. AI-based systems can identify pollinators from images, while drones and sensors track pollinator activity in real time (Koh *et al*., 2016). Mobile applications enable citizen scientists to contribute data, expanding monitoring networks globally (Silvertown, 2009). These tools improve early detection of declines and guide targeted interventions.

Community-based Approaches

Local communities play a central role in conservation. Farmer-led initiatives, such as planting pollinator-friendly crops or reducing pesticide use, have




Fig. 3 : Integrated framework of conservation strategies for pollinator recovery.

Table 2 : Conservation strategies and supporting evidence.

Strategy	Core approach	References
Habitat restoration	Wildflower strips, hedgerows, buffer zones, urban pollinator gardens	Kremen *et al.* (2007), Carvell *et al.* (2007), Baldock *et al.* (2015)
Pollinator-friendly farming	Reduced tillage, crop rotation, intercropping, organic and IPM systems	Garibaldi *et al.* (2014), Hole *et al.* (2005), Goulson (2013)
Policy frameworks	Global and regional pollinator initiatives, subsidies, certification schemes	IPBES (2016), Klein *et al.* (2007)
Genetic approaches	Disease-resistant honeybee breeding, genomic diversity conservation	Evans and Schwarz (2011), Zayed (2009), Cameron *et al.* (2011)
Technological monitoring	AI-based identification, drones and sensors, citizen-science applications	Koh *et al.* (2016), Silvertown (2009)
Community-based approaches	Farmer-led initiatives, education and outreach programs	Garibaldi *et al.* (2014), Kremen and Merenlender (2018)




Fig. 4 : Timeline of milestones in pollinator decline and conservation response.

proven effective (Garibaldi *et al*., 2014). Education and outreach programs raise awareness about pollinator importance and encourage behavioral change. Collaborative approaches that involve farmers, researchers, policymakers and citizens ensure long-term success (Kremen and Merenlender, 2018).

Future Directions

Future research and management efforts should prioritize landscape-scale, long-term monitoring networks that integrate satellite remote sensing, drone-based surveys and citizen-science data to detect early warning signs of decline across regions rather than isolated sites. Climate-resilient habitat design, including the strategic placement of floral corridors that anticipate shifting phenology, will be essential to maintain plant-pollinator synchrony under continued warming. Greater investment in participatory, community-led conservation models is needed to translate policy frameworks into on-ground practice, particularly in smallholder agricultural systems where top-down regulation has limited reach. Finally, interdisciplinary collaboration linking genomics, ecology, agronomy, and data science will be critical to developing predictive tools that can anticipate synergistic stressor interactions before they cause irreversible population collapse.

Conclusion

Pollinators are indispensable to agriculture, biodiversity, and ecosystem resilience. Their decline, driven by habitat loss, pesticides, pathogens, climate change and invasive species, poses serious risks to food security and ecological stability. Conservation strategies must therefore be comprehensive, integrating habitat restoration, pollinator-friendly farming, policy frameworks, genetic approaches, technological monitoring and community engagement. The evidence is clear that pollinator conservation is not only an ecological necessity but also an economic and social imperative. Protecting pollinators ensures sustainable crop production, supports rural livelihoods, and maintains biodiversity. Future efforts must emphasize global collaboration, combining scientific innovation with grassroots action, so that the foundation of ecosystems and the resilience of human societies are safeguarded.

References

Aizen, M.A. and Harder L.D. (2009). The global stock of pollinators. *Annu. Rev. Ecol., Evol. Systematics*, **40**, 25–45.

Baldock, K.C.R., Goddard M.A., Hicks D.M., Kunin W.E., Mitschunas N., Osgathorpe L.M. and Memmott J. (2015). Where is the UK's pollinator biodiversity? The importance of urban areas for flower-visiting insects. *Biological Conservation*, **192**, 448–457.

Biesmeijer, J.C., Roberts S.P.M., Reemer M., Ohlemuller R., Edwards M., Peeters T. and Kunin W.E. (2006). Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science*, **313(5785)**, 351–354.

Cameron, S.A., Lozier J.D., Strange J.P., Koch J.B., Cordes N.,

Solter L.F. and Griswold T.L. (2011). Patterns of widespread decline in North American bumble bees. *Proc. Nat. Acad. Sci.*, **108(2)**, 662–667.

Carvell, C., Meek W.R., Pywell R.F., Goulson D. and Nowakowski M. (2007). Comparing the efficacy of agri-environment schemes to enhance bumble bee abundance and diversity on arable field margins. *J. Appl. Ecol.*, **44(1)**, 29–40.

Evans, J.D. and Schwarz R.S. (2011). Bees brought to their knees: Microbes affecting honey bee health. *Trends Microbiol.*, **19(12)**, 614–620.

Gallai, N., Salles J.M., Settele J. and Vaissiere B.E. (2009). Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics*, **68(3)**, 810–821.

Garibaldi, L.A., Carvalheiro L.G., Leonhardt S.D., Aizen M.A., Blaauw B.R., Isaacs R. and Winfree R. (2014). From research to action: Enhancing crop yield through wild pollinators. *Front. Ecol. Environ.*, **12(8)**, 439–447.

Genersch, E. (2010). Honey bee pathology: Current threats to honey bees and beekeeping. *Appl. Microbiol. Biotechnol.*, **87(1)**, 87–97.

Goulson, D. (2013). An overview of the environmental risks posed by neonicotinoid insecticides. *J. Appl. Ecol.*, **50(4)**, 977–987.

Graystock, P., Yates K., Evison S.E.F., Darvill B., Goulson D. and Hughes W.O.H. (2013). The Trojan hives: Pollinator pathogens, imported and distributed in bumblebee colonies. *J. Appl. Ecol.*, **50(5)**, 1207–1215.

Hole, D.G., Perkins A.J., Wilson J.D., Alexander I.H., Grice P.V. and Evans A.D. (2005). Does organic farming benefit biodiversity? *Biological Conservation*, **122(1)**, 113–130.

Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services [IPBES] (2016). *The assessment report on pollinators, pollination and food production*. Bonn, Germany: IPBES Secretariat.

Kerr, J.T., Pindar A., Galpern P., Packer L., Potts S.G., Roberts S.M. and Pantoja A. (2015). Climate change impacts on bumblebees converge across continents. *Science*, **349(6244)**, 177–180.

Klein, A.M., Vaissiere B.E., Cane J.H., Steffan-Dewenter I., Cunningham S.A., Kremen C. and Tscharntke T. (2007). Importance of pollinators in changing landscapes for world crops. *Proc. Royal Soc. B*, **274(1608)**, 303–313.

Koh, I., Lonsdorf E.V., Williams N.M., Brittain C., Isaacs R., Gibbs J. and Ricketts T.H. (2016). Modeling the status, trends, and impacts of wild bee abundance in the United States. *Proc. Nat. Acad. Sci.*, **113(1)**, 140–145.

Kremen, C., Williams N.M., Aizen M.A., Gemmill-Herren B., LeBuhn G., Minckley R. and Ricketts T.H. (2007). Pollination and other ecosystem services produced by mobile organisms: A conceptual framework for the effects of land-use change. *Ecol. Lett.*, **10(4)**, 299–314.

Kremen, C. and Merenlender A.M. (2018). Landscapes that work for biodiversity and people. *Science*, **362(6412)**, eaau6020.

Memmott, J., Waser N.M. and Price M.V. (2004). Tolerance of pollination networks to species extinctions. *Proc. Royal Soc. B*, **271(1557)**, 2605–2611.

Michener, C.D. (2007). *The Bees of the World* (2nd ed.). Baltimore, MD: Johns Hopkins University Press.

Monceau, K., Bonnard O. and Thiery D. (2014). *Vespa velutina*: A new invasive predator of honeybees in Europe. *J. Pest Sci.*, **87(1)**, 1–16.

Morales, C.L. and Traveset A. (2009). A meta-analysis of impacts of alien vs. native plants on pollinator visitation and reproductive success of co-flowering native plants. *Ecol. Lett.*, **12(7)**, 716–728.

Ollerton, J., Winfree R., and Tarrant S. (2011). How many flowering plants are pollinated by animals? *Oikos*, **120(3)**, 321–326.

Pettis, J.S., Lichtenberg E.M., Andree M., Stitzinger J., Rose R. and vanEngelsdorp D. (2013). Crop pollination exposes honey bees to pesticides which alters their susceptibility to the gut pathogen *Nosema ceranae*. *PLoS ONE*, **8(7)**, e70182.

Potts, S.G., Biesmeijer J.C., Kremen C., Neumann P., Schweiger O. and Kunin W.E. (2010). Global pollinator declines: Trends, impacts and drivers. *Trends Ecol. Evol.*, **25(6)**, 345–353.

Rosenkranz, P., Aumeier P. and Ziegelmann B. (2010). Biology and control of *Varroa destructor*. *J. Invertebrate Pathol.*, **103(Suppl. 1)**, S96–S119.

Rundlof, M., Andersson G.K.S., Bommarco R., Fries I., Hederstrom V., Herbertsson L. and Smith H.G. (2015). Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature*, **521(7550)**, 77–80.

Silvertown, J. (2009). A new dawn for citizen science. *Trends Ecol. Evol.*, **24(9)**, 467–471.

Steffan-Dewenter, I. and Tscharntke T. (1999). Effects of habitat isolation on pollinator communities and seed set. *Oecologia*, **121(3)**, 432–440.

Stout, J.C. and Morales C.L. (2009). Ecological impacts of invasive alien species on bees. *Apidologie*, **40(3)**, 388–409.

van Engelsdorp, D., Hayes J., Underwood R.M. and Pettis J. (2009). A survey of honey bee colony losses in the United States, fall 2008 to spring 2009. *J. Apicult. Res.*, **48(1)**, 7–14.

Whitehorn, P.R., O'Connor S., Wackers F.L. and Goulson D. (2012). Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science*, **336(6079)**, 351–352.

Zayed, A. (2009). Bee genetics and conservation. *Apidologie*, **40(3)**, 237–262.