



Plant Archives

Journal homepage: <http://www.plantarchives.org>

DOI Url : <https://doi.org/10.51470/PLANTARCHIVES.2025.v25.no.1.252>

EVALUATION OF BIO-EFFICACY OF *BACILLUS SUBTILIS* BASED BIO-FUNGICIDE (TAEGRO) AGAINST POWDERY MILDEW DISEASE OF GRAPES

Ramesh^{1*}, Manjunath Hubballi² and Raghavendra Achari³

¹Main Horticulture Research and Extension Centre, UHS, Bagalkot - 587 104, Karnataka, India

²Directorate of Research, UHS, Bagalkot - 587 104, Karnataka, India

³College of Horticulture, Munirabad-583 233, Karnataka, India

*Corresponding author E-mail: ramesh3963@gmail.com

(Date of Receiving-19-02-2025; Date of Acceptance-28-04-2025)

ABSTRACT

Grape (*Vitis vinifera* L.) is an important cash crop in India and powdery mildew is a major biotic disease which causes severe loss in yield and quality of grapes. An field experiments was conducted during the fruiting season of October–March 2016-17 to study the efficacy of *Bacillus subtilis* based bio fungicides (Taegro) against powdery mildew diseases of grapes. Alone eight sprays of triadimefon 75% WP at 1g/liter recorded lowest powdery mildew disease 7.47 PDI. First five sprays of triadimefon 75% WP at 1g/liter followed by three sprays of *Bacillus subtilis* based bio fungicides (Taegro) at 0.370g/liter and alone first five sprays of Triadimefon 75% WP at 1g/liter reduced the PDI of 11.43 and 12.27 respectively. Alone eight sprays of *Bacillus subtilis* based bio fungicides (Taegro) at 0.185, 0.370 and 0.50 g/liter recorded the PDI of 25.03, 22.93 and 22.43 respectively. Under low to moderate disease pressure conditions, the efficacy of Triadimefon 75% WP at 1g/liter and *Bacillus subtilis* based bio fungicides (Taegro) at 0.370g/liter in controlling powdery mildew was on par with alone spray of Triadimefon 75% WP at 1g/liter. The management of powdery mildew disease by using fungicides which increases the input cost and affect berry quality. Under high disease pressure condition, *Bacillus subtilis* based bio fungicides (Taegro) were not so effective when used alone, but was effective when used in integration with fungicide sprays.

Key words : Grapes, *Bacillus subtilis*, Triadimefon 75% WP, Powdery mildew

Introduction

Grape (*Vitis vinifera* L.) is one of the finest fruits and the healthiest food and it is considered one of the most popular and common fruits in the world. Taxonomically, grapes belong to the family Vitaceae which is classified into two sub-genera, Euvitis (2n=38) Planch and Muscadinia (2n=40) Planch (Einset and Pratt, 1975). Most cultivated grape varieties belong to *Vitis vinifera* L., which originated in Eurasia and spread by man all the way through the world. Grapes fruits are rich source of minerals like calcium and iron and vitamins A, C and B1 and B2 that can contribute to a balanced healthy life. In addition, it has commendable medicinal qualities and has been used in neurotherapy for centuries. The flavonoids present in grapes act as antioxidants and reduce the damage caused by free radicals. Due to its medicinal

properties, grape juice is popularly known as ‘nectar of the gods’. The grape was introduced to India in 1300 AD by Muslim invaders from Iran and Afghanistan (Thapar, 1960). Grapevine is grown in over 80 countries across 7.2 million hectares, producing approximately 74.7 million tonnes annually (OIV, 2023). In India, grape is cultivated on about 1.58 lakh hectares, with an annual production of 3.4 million tonnes and a productivity of 21.5 tonnes/ha (NHB, 2023).

The major threat in grape production is powdery mildew (*Erysiphe necator* Schw) disease and it affects all green succulent tissues. The disease is quite severe in India and foliar infections reduce the photosynthetic activity of grapevines (Chadha and Shikhamany, 1999), while fruit infections impair wine characteristics or reduce the yield and market value of table grapes (Sawant and

Sawant, 2008). Infections on the pedicle lead to shriveling of the clusters during storage reducing the shelf life (Gubler, *et al.*, 2013). The disease is favored by warm, humid and cloudy weather (Pearson, 1988).

Material and methods

A field experiment was conducted three years old vein orchard at Horticulture Research and Extension Station, Vijayapur (Tidagundi) to study the bio-efficacy of *Bacillus subtilis* based bio fungicides (Taegro) against powdery mildew diseases of grapes. It was conducted in randomized block design with ten treatments (consisting of alone or alternative sprays of *Bacillus subtilis* bio fungicides/triadimefon 75% WP) with three replications at sprayed weekly interval during 2016-17. The vines were supplied with organic fertilizer in the form of vermin-compost and inorganic chemical fertilizer as per recommendation mentioned in package of practices for Horticultural crops, UHS, Bagalkot. Three rows for each replication with five plants in a row were selected. The three plants in middle row excluding the terminal vines on each side were selected for the recording observation. The treatments were initiated at the first disease appearance stage (DAS) of powdery mildew disease. The required quantity of fungicide/ bio-agent as per treatments were measured out/weighed with measuring cylinder/balance and suitable dissolved in a required quantity of water to obtain spray solution of desired concentrations. Observations were recorded at before first spray and later at 5 days after 3rd, 5th and 8th spray. The disease incidence on leaves and bunches were recorded by using 0-4 scale (Horsfall and Heuberger, 1942).

Disease rating/ Grade	Per cent area (leaves/ bunch) covered/infected
0	No disease/ infection
1	<25% area covered/infected
2	26-50% area covered/infected
3	51-75% area covered/infected
4	>75% area covered/infected

Further, the scored data were converted into Per cent Disease Index (PDI) of plants using formula given by McKinney (1923).

$$\text{Per cent Disease Index (PDI)} = \frac{\text{Sum of numerical rating}}{\text{Number of bunches or leaves observed}} \times \frac{100}{\text{Max. rating}}$$

The data was suitably transformed and analyzed statistically. The marketable fruit yield in terms of kilograms per vine and later converted to tonner per hectare and analyzed statistically.

Results and Discussion

Alone sprays of *Bacillus subtilis* based bio fungicides (Taegro) at 0.185, 0.370 and 0.50 g/liter when applied as eight weekly sprays reduced the progress of powdery mildew disease by 26.32%, 32.50% and 33.97% with a PDI of 25.03, 22.93 and 22.43 respectively when compared to untreated control (33.97 PDI) at five days after 8th spray. Alone eight sprays of triadimefon 75% WP at 1g/liter reduced the progress of powdery mildew disease by 78.01% with a PDI of 7.47 per cent. First five sprays of triadimefon 75% WP at 1g/liter followed by three sprays of *Bacillus subtilis* based bio fungicides (Taegro) at 0.370g/liter and alone first five sprays of triadimefon 75% WP at 1g/liter reduced the progress of powdery mildew disease by 66.35% and 63.88% with a PDI of 11.43 and 12.27 respectively. Both these two treatments were statistically on par with each other (Table 1).

Alone eight sprays of triadimefon 75% WP at 1g/liter recorded highest productivity of 13.25 tons/ha which was on par with treatment first five sprays of triadimefon 75% WP at 1g/liter followed by three sprays of *Bacillus subtilis* based bio fungicides (Taegro) at 0.370g/liter (11.90 tons/ha). However, the treatment triadimefon 75% WP at 1g/liter followed by three sprays of *Bacillus subtilis* based bio fungicides (Taegro) at 0.370g/liter was statistically on par with five sprays of triadimefon 75% WP at 1g/liter (10.55 tons/ha). Alone eight sprays of *Bacillus subtilis* based bio fungicides (Taegro) at 0.185, 0.370 and 0.50 g/liter concentration recorded productivity of 4.96, 6.12 and 7.72 tons/ha respectively. The yields in these treatments were increased from 2.04, 3.03 and 3.82 times more than the untreated control. Increasing yield over untreated control was achieved 6.56, 5.69 and 5.17 times in treatments. Alone eight sprays of triadimefon 75% WP at 1g/liter, five sprays of triadimefon 75% WP at 1g/liter followed by three sprays of *Bacillus subtilis* based bio fungicides (Taegro) at 0.370g/liter and alone five sprays of triadimefon 75% WP at 1g/liter respectively (Table 1).

This is in agreement with the findings Singh, P.N. (2017) reported that bio-agents *Ampelomyces quisqualis*, *Trichoderma harzianum* and *Saccharomyces cerevisiae* can be used as alternative to chemicals fungicides for control of powdery mildew disease of grapes. DR-38, DR-39, TL-171, DRo-198, TS-204, TS-205, and DR-219 strains of *Bacillus sp* can reduce downy mildew, powdery mildew and anthracnose disease of grapes (Indu S. Sawant. *et al.*, 2016). Sharma,

Table 1: Bio-efficacy of *Bacillus subtilis* based bio-fungicide (*Taegro*) against powdery mildew disease of Grapes (2016-17)

Tr. No	Treatments	Formulation/dosage	Weekly application sequence	Powdery mildew Disease (PDI)				% Reduction over control after 8 th spray	Productivity (marketable fruit yield)		Increase in productivity (times than control)
				Before first spray	5 days after 3 rd spray	5 days after 5 th spray	5 days after 8 th spray		kg/plant	t/ha	
1	Control	--	--	2.23 (8.59)*	19.57 (26.26)	25.40 (30.26)	33.97 (35.65)		0.89	2.02 (8.17)	---
2	Triadimefon 75% WP	1.00 g/l	ABCDEFGH	2.23 (8.59)	8.07 (16.50)	9.13 (17.58)	7.47 (15.86)	78.01	5.24	13.25 (21.35)	6.56
3	<i>Bacillus subtilis</i> (commercial formulation)	2.00 ml/l	ABCDEFGH	2.23 (8.59)	15.20 (22.95)	23.37 (28.90)	28.47 (32.25)	16.19	1.62	4.14 (11.74)	2.04
4	<i>Taegro</i>	0.185 g/l	ABCDEFGH	2.23 (8.59)	14.80 (22.63)	15.67 (23.32)	25.03 (30.02)	26.32	1.45	4.96 (12.87)	2.46
5	<i>Taegro</i>	0.370 g/l	ABCDEFGH	2.23 (8.59)	13.53 (21.58)	14.50 (22.38)	22.93 (28.61)	32.50	2.99	6.12 (14.32)	3.03
6	<i>Taegro</i>	0.500 g/l	ABCDEFGH	2.23 (8.59)	11.97 (20.24)	12.80 (20.96)	22.43 (28.26)	33.97	3.10	7.72 (16.13)	3.82
7	Triadimefon 75% WP <i>Taegro</i>	1.00 g/l 0.370 g/l	ABCDE FGH	2.23 (8.59)	8.50 (16.95)	10.43 (18.84)	11.43 (19.76)	66.35	5.56	11.90 (19.99)	5.69
8	Triadimefon 75% WP	1.00 g/l	ABCDE	2.23 (8.59)	8.50 (16.95)	10.43 (18.84)	12.27 (20.50)	63.88	5.41	10.55 (18.96)	5.17
9	Triadimefon 75% WP <i>Taegro</i>	1.00 g/l 0.370 g/l	ACE BDFGH	2.23 (8.59)	8.27 (16.71)	11.63 (19.93)	17.67 (24.86)	47.98	3.42	8.86 (17.31)	4.39
10	Triadimefon 75% WP	1.00 g/l	ACE	2.23 (8.59)	8.27 (16.71)	13.37 (21.45)	19.47 (26.18)	42.68	3.29	7.58 (15.98)	3.75
	SEM±			NS	0.24	0.38	0.51			0.47	
	CDat5%			NS	0.70	1.14	1.53			1.40	
	CV%			NS	3.54	4.51	4.42			25.55	

* Figures in parenthesis are angular transformed value

A. K.(2017) reported that application of chlorine dioxide (AgroRakshak) at 150 ppm was found effective in controlling powdery mildew of grapes. Pre harvest treatment of grapes with *T. harzianum*5R will increase fruit freshness and overcome many postharvest diseases (S.D. Sawant and Indu S. Sawant. 2010).

References

- A.K. Sharma., Indu, S. Sawant., S. D. Sawant., Sujoy saha., Pratiksha Kadam and R. G Somkumar. (2017). Aqueous chlorine dioxide for the management of powdery mildew vis-à-vis maintaining quality of grapes and raisins. *J. Eco-friendly agriculture*. **12(2)**: 59-64.
- Chadha, K. L. and Shikhamany, S.D. (1999) The Grape-improvement, Production and Postharvest Management. Malhotra Publishing House, New Delhi.
- Einset, J. and C. Pratt, (1975). Grapes. In: J. Janick and J.N. Moore (Eds.), *Advances in Fruit Breeding*, pp. 130–153. Purdue University Press, West Lafayette.
- Gubler, W.D., Hashim, J.M., Smilanick, J.L. and Leavitt, G.M.(2013). Post harvest diseases of table grapes. In: Bettiga, L.J. (Ed.), *Grape Pest Management*, third ed. University of California, California, 133–136.
- Horsfall, J. G. and Heuberger, J. W. (1942). Measuring magnitude of a defoliation disease in tomatoes. *Phytopathology*, **32**: 226–232.
- Indu, S. Sawant., Pallavi, N. Wadkar., Yogita, R. Rajguru., Nileema, H. Mhaske., Varsha P. Salunkhe., S. D. Sawant and Anuradha Upadhyay. (2016). Biocontrol potential of two novel grapevine associated *Bacillus* strains for management of anthracnose disease caused by *Colletotrichum gloeosporioides*. *Biocontrol Science and Technology*., **26(7)**: 964–979.
- McKinney, H. H. (1923). A new system of grading plant diseases. *Journal of Agriculture Research*, **26**: 195–218.
- National Horticultural Board. (2023). Horticultural statistics at a glance. Ministry of Agriculture Farmers Welfare, Government of India. <https://www.nhb.gov.in>
- OIV. (2023). *Statistical Report on World Vitiviniculture*. <https://www.oiv.int>
- Pearson, R.C. (1988). Powdery mildew. In: Pearson, R.C., Goheen, A.C. (Eds.), *Compendium of Grape Diseases*. The American Phytopathological Society, APS Press, 9–11.
- S.D. Sawant and Indu, S. Sawant. (2010). Improving the shelf life of grapes by pre-harvest treatment with *Trichoderma*

- harzianum* 5R. *J. Eco-friendly Agriculture*, **5(2)**: 179-182.
- Sawant, S.D. and Sawant, I.S. (2008). Use of potassium bicarbonate for the control of powdery mildew in table grapes. *Acta Hort.* **785**: 285–291.
- Singh, P. N., Singh, S.K., Tetali, S. P. and Lagashetti, A.C. (2017). Biocontrol of powdery mildew of grapes using culture filtrate and biomass of fungal isolates. *Plant Pathology & Quarantine*, **7(2)**: 181–189.
- Thapar, A.R. (1960). Horticulture in the hill regions of North India. Directorate of Extension, Ministry of Food and Agriculture, New Delhi.